

Effective widths and integrated cross sections for two-photon transitions in ^{178}Hf

Silviu Olariu and Agata Olariu

Institute of Physics and Nuclear Engineering,
Department of Fundamental Experimental Physics
76900 Magurele, P.O. Box MG-6, Bucharest, Romania
e-mail: olariu@roifa.ifa.ro

23 February 1999

The possibility to produce an increase of the activity of an isomeric sample by irradiating the sample with a laser pulse and the x-ray pumping of nuclear transitions have been discussed some time ago. [1] We have studied recently the rates for two-photon transitions from isomeric states to given final states, induced by an incident flux of photons having a continuous energy distribution. [2],[3]

In this work we have calculated integrated cross sections for two-photon processes in ^{178}Hf . The calculations are based on tabulated nuclear data. [4] We have assumed that the nucleus is initially in a state $|i\rangle$ of energy E_i and spin J_i , which in general is different from the isomeric state. By absorbing an incident photon of energy E_{ni} , the nucleus makes a transition to a higher intermediate state $|n\rangle$ of energy E_n , spin J_n and half-life t_n . The state $|n\rangle$ then decays into a lower state $|l\rangle$ of energy E_l and spin J_l by the emission of a gamma-ray photon having the energy E_{nl} or by internal conversion. In some cases the state $|l\rangle$ may be situated above the initial state $|i\rangle$, and in these cases the transition $|n\rangle \rightarrow |l\rangle$ is followed by further gamma-ray transitions to lower states.

We have analyzed two-photon transitions in ^{178}Hf for which there is an intermediate state $|n\rangle$ of known energy E_n , spin J_n and half-life t_n , for which the intermediate state is connected by a known gamma-ray transition to the initial state $|i\rangle$ and to a lower state $|l\rangle$, and for which the relative intensities R_{ni} , R_{nl} , $R_{nl'}$ of the transitions to lower states are known.

The cross sections are summed over all lower states $|l\rangle$ which are possible for a given pair of initial and intermediate states $|i\rangle$, $|n\rangle$. The integrated cross sections

are calculated as

$$\sigma_{int}^{tot} = \frac{2J_n + 1}{2J_i + 1} \frac{\pi^2 c^2 \hbar^2}{E_{ni}^2} \hbar \Gamma_{eff}^{tot}, \quad (1)$$

where

$$\Gamma_{eff}^{tot} = \sum_{l \neq i} \Gamma_{eff}, \quad (2)$$

and where $E_{ni} = E_n - E_i$. The quantity Γ_{eff} is the effective width of the two-photon transition,

$$\Gamma_{eff} = F_R \ln 2 / t_n, \quad (3)$$

where t_n is the half-life of the intermediate state $|n\rangle$, and the dimensionless quantity F_R has the expression

$$F_R = \frac{(1 + \alpha_{nl}) R_{ni} R_{nl}}{[(1 + \alpha_{ni}) R_{ni} + (1 + \alpha_{nl}) R_{nl} + \sum_{l'} (1 + \alpha_{nl'}) R_{nl'}]^2}. \quad (4)$$

In Eq. (4), $\alpha_{ni}, \alpha_{nl}, \alpha_{nl'}$ are the internal conversion coefficients for the transitions $|n\rangle \rightarrow |i\rangle, |n\rangle \rightarrow |l\rangle, |n\rangle \rightarrow |l'\rangle, l' \neq i, l$.

The effective widths Γ_{eff}^{tot} and the integrated cross sections σ_{int}^{tot} are given in Table I. The multiplicities of the transitions are given in columns ni and nl . The largest integrated cross section found among the listed 24 two-photon processes in ^{178}Hf has the value $1.65 \times 10^{-26} \text{ cm}^2 \text{ keV}$. If the initial state is the isomeric state at $E_i=2446 \text{ keV}$, the nuclear parameters required for the calculation of the two-photon integrated cross section are known at present only for the intermediate state at $E_n=2573 \text{ keV}$.

References

- [1] S. Olariu, *Essay on the gamma ray laser*, private communication to C. B. Collins, 1979; S. Olariu, I. I. Popescu and C. B. Collins, Rev. Roum. Phys. **27**, 559 (1982).
- [2] S. Olariu and A. Olariu, Phys. Rev. C **58**, 333 (1998).
- [3] S. Olariu and A. Olariu, Phys. Rev. C **58**, 2560 (1998).
- [4] *Table of Isotopes*, 8th ed., edited by R. Firestone (Wiley, New York, 1996); National Nuclear Data Center Online Service, Brookhaven National Laboratory, New York; F. Rösler, H. M. Fries, K. Alder and H. C. Pauli, Atomic Data and Nuclear Data Tables **21**, 291 (1978).

Table 1: Effective width Γ_{eff}^{tot} and integrated cross section σ_{int}^{tot} for two-photon transitions in ^{178}Hf .

E_i (keV)	E_n (keV)	ni	nl	Γ_{eff}^{tot} (eV)	σ_{int}^{tot} (cm ² keV)
1434.2	1496.4	E2	E2,M1	3.32×10^{-6}	1.65×10^{-26}
0	1174.6	E2	E2	1.80×10^{-4}	2.51×10^{-27}
0	1496.4	E2	M1,E2	9.97×10^{-5}	8.56×10^{-28}
93.2	1174.6	E2	E2	1.79×10^{-4}	5.89×10^{-28}
93.2	1496.4	M1	E2	1.26×10^{-4}	2.46×10^{-28}
0	1276.7	E2	M1,E2	8.26×10^{-6}	9.73×10^{-29}
93.2	1276.7	M1	E2	1.63×10^{-5}	4.47×10^{-29}
1513.8	1636.7	E1	E1,M1,E2	1.29×10^{-7}	4.00×10^{-29}
1554.0	1636.7	E1	E1,M1,E2	7.57×10^{-8}	3.60×10^{-29}
306.6	1276.7	E2	M1,E2	1.18×10^{-5}	2.67×10^{-29}
306.6	1496.4	E2	M1,E2	1.26×10^{-5}	1.90×10^{-29}
306.6	1174.6	E2	E2	6.06×10^{-6}	1.72×10^{-29}
1538.8	1636.7	M1	E1,M1,E2	1.69×10^{-8}	8.27×10^{-30}
1409.4	1636.7	E2	E1,M1,E2	7.05×10^{-8}	6.41×10^{-30}
1433.6	1636.7	E2	E1,M1,E2	3.31×10^{-8}	4.85×10^{-30}
1512.6	1636.7	M1	E1,M1,E2	7.12×10^{-9}	1.78×10^{-30}
306.6	1636.7	E1	E1,M1,E2	2.85×10^{-7}	7.56×10^{-31}
1384.5	1554.0	E2	E2,M2	2.36×10^{-10}	4.55×10^{-32}
632.2	1554.0	E2	E2,M2	1.41×10^{-9}	6.37×10^{-33}
306.6	1554.0	E2	E2,M2	1.25×10^{-9}	4.46×10^{-33}
1147.4	1554.0	M2	E2	1.35×10^{-10}	2.40×10^{-33}
2433.3	2573.5	M1	E2,M2	6.14×10^{-13}	1.29×10^{-34}
2446.1	2573.5	M2	M1,E2	9.86×10^{-14}	2.09×10^{-35}
2136.5	2573.5	E2	M1,M2	5.93×10^{-13}	1.38×10^{-35}